FOR USE IN ANIMALS ONLY

INDICATIONS: For use as an aid in controlling minor bleeding from superficial cuts and wounds and after dehorning.

DOSAGE AND ADMINISTRATION:

Apply powder freely to bleeding surface. Repeat as needed. Bandage if necessary.

CONTRAINDICATIONS: In case of deep or puncture wounds or serious burns, consult veterinarian. If redness, irritation, or swelling persists or increases, discontinue use and consult veterinarian.





OBSERVE LABEL DIRECTIONS

Manufactured for: DURVET, INC. Blue Springs, Missouri 64014

B-1101-07 Rev. 12-10

durvet

BLOOD STOP POWDER









Controls Minor Bleeding

www.durvet.com

NET CONTENTS: 16 oz (2.205 Kg)

KEEP OUT OF REACH OF CHILDREN

		100		44	
A	ctive	e Ing	gre	dıe	nts

Ferrous Sulfate • /H ₂ O	84%
nactive Ingredients:	
Ammonium Alum	. 5%
Chloroxylenol	. 1%
Tannic Acid	. 1%
Free flowing absorbent base	. 9%
(Non-Sterilized)	

For external use only.

Store at controlled room temperature between 15° and 30°C (59°–86°F)

Keep container tightly closed.

LOT NO.:

EXP. DATE:

