ferti-lome

Triple Action

Insecticide • Fungicide • Miticide

• Can Be Used Up To Day of Harvest

 For Control of Insects, Diseases, and Mites on Vegetables, Fruits, Nuts, Herbs, Spices, Roses, Flowers and Shrubs Around the Home

KEEP OUT OF REACH OF CHILDREN

CAUTION

ACTIVE INGREDIENTS:

Pyrethrins	0.25%
Clarified Hydrophobic Extract of Neem Oil.	70.00%
OTHER INGREDIENTS:	<u>29.75%</u>
TOTAL:	100.00%

CONTAINS NEEM

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS ONE PINT (32 FL. OZS. / 946 ML)

lf On Skin Or Clothing	FIRST AID Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
lf Swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
lf Inhaled	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
lf In Eyes	Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. Call a poison control center or doctor for treatment advice.
	uct container or label with you when calling a poison control center ping for treatment.

fertielome[®] Triple Action

7

32221

12246118



PEEL BACK BOOK HERE

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic organisms. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

BEE HAZARD

This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION: For organic gardening.

SHAKE WELL BEFORE AND DURING USE.

Multi-purpose insecticide/fungicide/miticide. Kills eggs, larval and adult stages of listed insects. Prevents fungal attack of plants' tissues. Not for use in food handling areas. **ferti**•lome®

Triple Action is most effective when applied in early to midmorning or late afternoon when adult insect pests, such as whiteflies, are normally sedentary on the undersides of the foliage. Use spray solutions within several hours of preparation for maximum effectiveness. Do not use more than 1 time per day on the same plants. Do not apply this product more than 10 times per season. For best results maintain constant agitation in spray equipment. For optimal performance, do not mix with cold water or apply under cold ambient temperatures (less than 45°F). If product thickens (freezes) before using, place in a warm room (greater than 45°F) for a few days or in warm water (120°F) for 2 to 8 hours until completely thawed. Do not use partially thawed material. Do not apply to wilted or otherwise stressed plants, or to newly transplanted material prior to root establishment. When using in conjunction with beneficial insects, conduct a small trial to assure compatibility before using on a large scale. As with other products that contain oils, exercise care in timing application to early morning/late evening to minimize the potential for leaf burn. Do not add surfactants (spreaders, stickers, extenders, etc.) to ferti•lome® Triple Action

Use with care on plants with tender tissue. Do not apply to known spray sensitive plants species such as: Impatiens flowers, Fuchsia flowers, Hibiscus flowers, some Rose flowers, ornamental Olive Trees, Red Maples, some Carnation varieties or similar plants without prior testing. Not for use in outdoor residential misting systems. Do not allow adults, children or pets to enter the treated area until sprays have dried. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) or apply when heavy rain is expected. Rinse applicator over lawn or garden areas only.

FUNGICIDE USE

For use on FRUITS, NUTS, VEGETABLES, HERBS, SPICES, ROSES, HOUSEPLANTS, FLOWERS and SHRUBS.

Application Instructions: This product is an effective fungicide for the prevention and control of various fungal diseases including Powdery Mildew, Black Spot, Brown Spot, Dollar Spot. Snow Mold. Downy Mildew. Anthrachose. Rust. Leaf Spot. Botrytis, Needle Rust, Scab and Flower, Twig and Tip Blight. As a preventative, apply fertiolome® Triple Action on a 7 to 14 day schedule until the potential for disease development is no longer present. To control disease already present, apply this product on a 7 day schedule until disease pressure is eliminated. Then continue spraving on a 14 day schedule to prevent the disease from reoccurring. To prevent Rust Leaf Spot diseases, Anthracnose and Scab, begin applications at the first sign of spring budding. To prevent Powdery Mildew, apply in midsummer or when disease is first detected. For Brown Patch and Dollar Spot, begin spraying at first sign of disease on a 7 to 14 dav interval.

For Snow Mold, begin spraying on a 14 day interval in the fall before the first snow. Continue applications until disease pressure no longer exists.

INSECTICIDE USE

For use on FRUITS, NUTS, VEGETABLES, HERBS, SPICES, ROSES, HOUSEPLANTS, FLOWERS and SHRUBS.

Application Instructions: Apply **ferti+lome® Triple Action** at first sign of insects/mites. This product provides for a quicker and more complete control, and acts as an "exciter" to flush insect pests out of hiding and into direct contact with spray residues. For control of Aphids, Spider Mites, Scale, Mealybugs, Whiteflies, Beetles, Loopers, Leafminers, Leafrollers, Armyworms, Webworms, Weevils, Tent Caterpillars, Fungus Gnats, Leafhoppers, Cabbageworms, Squash Bugs, Squash Vine Borer, Stink Bugs, Thrips, Tomato Hornworm, Lace Bugs, Boxelder Bugs, Grasshoppers, Crickets, Ants, Earwigs, Chinch Bugs, Chiggers, and many other insect pests. This product is most effective when applied on a 7 to 14 day schedule. For heavy insect populations, spray on a 7 day schedule.

Mixing Instructions: Mix **ferti-lome® Triple Action** at the rate of 2 tablespoons (1 fluid ounce) per gallon of water. Thoroughly mix solution and spray all plant leaf surfaces (including undersides of leaves) until completely wet. Excessive application is unnecessary and to be avoided. Frequently agitate solution as you spray.

STORAGE AND DISPOSAL

Pesticide Storage: Do not store diluted solutions for later use.

Pesticide Disposal and Container Handling: Nonrefillable container. Do not reuse or refill container. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY

To the extent consistent with applicable law, Voluntary Purchasing Groups, Inc. (VPG) warrants that the material contained herein conforms to the description on the label and is reasonable fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the insect problem, condition of the crop, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are all beyond the control of the seller. To the extent consistent with applicable law, Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OF MERCHANTABILITY IS MADE. Manufactured By:



230 FM 87 • BONHAM, TX 75418 EPA Reg. No. 70051-111-7401 EPA Est. No. 7401-TX-01 Visit Us At: www.fertilome.com Product Questions? 855-270-4776

12246-0916-CL

