

ferti·lome®



Weed Free Zone PLUS LAWN FERTILIZER 18-0-6


- Kills Over 200 Broadleaf Weeds As Listed
- Controls Tough Broadleaf Weeds In Established Lawns
- Superior Broadleaf Weed Control In Cool Weather

ACTIVE INGREDIENT:	
2,4-D, 2-ethylhexyl ester	1.153%
Mecoprop-p acid0174%
Dicamba acid073%
Sulfentrazone034%
OTHER INGREDIENTS:	98.566%
TOTAL	100.000%
THIS PRODUCT CONTAINS:	
15.30 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.765%. CAS 1928-43-4.	
3.48 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.174%. CAS 16484-77-8.	
1.46 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.073%. CAS 1918-00-9.	
0.68 lb N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide per ton or 0.034%. CAS 122836-35-5.	
Isomer Specific by AOAC Methods	
TRIMEC® is a registered trademark of PBI/Gordon Corporation.	

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Back Panel for Additional Precautionary Statements, First Aid and Directions for Use.

NET WEIGHT 19 LBS. (8.62 KG)



**Weed Free Zone
Plus Lawn Fertilizer**

10930

Weed Free Zone Plus Lawn Fertilizer

Weed Free Zone Plus Lawn Fertilizer

Covers Up To
7,600 Sq. Ft.

Weed Free Zone PLUS LAWN FERTILIZER

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing. Wear long-sleeved shirt and long pants, shoes plus socks and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

FIRST AID

IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-858-7378 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For Residential Use Only.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

INSTRUCTIONS

ferti•lome® Weed Free Zone Plus Lawn Fertilizer is a combination of four proven weed killers – 2,4-D, mecoprop-p, dicamba and sulfentrazone. This herbicide controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 18-0-6 fertilizer provides a supply of plant nutrients.

USE PRECAUTIONS

Do not use on Dichondra, Carpetgrass, St. Augustinegrass, or Turfgrass where desirable clovers are present. Do not apply this product to newly seeded grasses until well-established or after the second or third mowing. Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area. This product will not prevent weed seeds from producing new plants.

Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

HOW MUCH TO APPLY

Apply 2.5 to 3.0 pounds of product per 1,000 square feet of turfgrass. A second application may be made in approximately 30 days.

Do not apply more than 3 pounds of product per 1,000 square feet of turfgrass per application. If necessary, a second application may be made no sooner than 30 days after the first. Do not apply more than 6 pounds of product per 1,000 square feet per year.

WHEN TO APPLY

Apply when broadleaf weeds are actively growing in spring or early fall.

- Apply when the air is calm (choose a calm day) to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Possible retreatment may be necessary if lawn is heavily overrun by weeds, if adverse low moisture conditions prevail, or if weeds are in a state of poor growth.

WHAT TO EXPECT

- Mow lawn to normal height 1 to 2 days before application.
- Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
- Apply when broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew, rainfall or irrigation/ watering aids in control. If grass is not moist at time of application, sprinkle lightly with water.
- Avoid applying when temperatures are consistently above 90°F.
- Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

DISTRIBUTION AND SPREADER PATTERNS

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

Coverage – To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to 7,600 square feet.

APPLICATION RATE, LBS./1,000 SQ. FT.	TREATED AREA FOR THIS BAG SIZE, SQ. FT.
2.50	7,600
2.75	6,900
3.00	6,300

SINGLE PASS DISTRIBUTION

For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps.

- For rotary spreaders, operate at intervals corresponding to the swath width. For drop spreaders, overlap the wheel tracks to ensure complete coverage.
- First, spread the borders of your lawn and header strips for turning around. Then apply this product to the center with straight line passes
- Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

CALIBRATION INSTRUCTIONS AND SPREADER SETTINGS

Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended rate under actual application conditions.

To provide proper distribution, calibrate spreader before application.

- For 2.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 400 square feet and adjust to apply at the rate of 1 pound of product per 400 square feet.
- For 2.75 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 360 square feet and adjust to apply at the rate of 1 pound of product per 360 square feet.
- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.

10930

SPREADER GUIDE / ONE BAG WILL COVER UP TO 7,600 SQ. FT.

SPREADER	SPREADER SETTINGS Lbs. of Product/1000 Sq. Ft.		
	2.5 LBS.	2.75 LBS.	3.00 LBS.
Scotts®/Republic Accugreen (Drop)	2 ¾	3	3 ¼
Scotts®/Republic Speedy Green (Broadcast)	3	3 ¼	3 ½
ferti•lome®/EarthWay Ev-N-Spred (Broadcast)	13	13 ½	14

These settings can be used only as a guide, they are not absolute. The speed at which the applicator walks, spreader condition and the texture of the turf will alter the rate of application.

BROADLEAF WEEDS

Annual Yellow Sweetclover, Aster, Austrian Fieldcress, Bedstraw, Beggarticks, Betony-Florida, Bindweed-Field, Bird Vetch, Bitter Wintercress, Bittercress-Hairy, Bitterweed, Black-Eyed Susan, Black Medic, Black Mustard, Blackseed Plantain, Blessed Thistle, Bloodflower Milkweed, Blue Lettuce, Blue Vervain, Bracted Plantain, Bristly Oxtongue, Broadleaf Dock, Broadleaf Plantain, Broomweed, Buckhorn, Buckhorn Plantain, Bulbous Buttercup, Bull Thistle, Bullnettle, Burclover, Burdock, Burning Nettle, Burweed, Buttercup, Buttonweed, Canada Thistle, Carolina Geranium, Carpetweed, Catchweed Bedstraw, Catnip, Catsear, Chickweed-Common, Chickweed-Mouseear, Chicory, Cinquefoil, Clover-Crimson, Clover-Hop, Clover-Red, Clover-Strawberry, Clover-Sweet, Clover-White, Cockle, Cocklebur, Common Mullein, Creeping Jenny, Cudweed, Curly Dock, Daisy-English, Daisy-Fleabane, Daisy-Oxeye, Dandelion, Dichondra, Dogfennel, Dollarweed, False Dandelion, Falseflax, False Sunflower, Fiddleneck, Florida Pusley, Frenchweed, Galinsoga, Goldenrod, Ground Ivy, Gumweed, Hairy Fleabane, Hawkweed, Healall, Heartleaf Drymary, Heathaster, Hedge Bindweed, Hedge Mustard, Henbit, Hoary Cress, Hoary Plantain, Hoary Vervain, Horsenettle, Jimsonweed, Knawel, Knotweed, Kochia, Lambsquarters, Lespedeza, Mallow, Matchweed, Mexicanweed, Milk Vetch, Morningglory, Mouseear Hawkweed, Mugwort, Musk Thistle, Mustard, Narrowleaf Plantain, Narrowleaf Vetch, Nettle, Orange Hawkweed, Oxalis, Parsley-Piert, Parsnip, Pearlwort, Pennygrass, Pepperweed, Pigweed, Pineywoods Bedstraw, Plains Coreopsis, Plantain, Poison Ivy, Poison Oak, Pokeweed, Poorjoe, Prairie Sunflower, Prickly Lettuce, Prickly Sida, Prostrate Knotweed, Prostrate Pigweed, Prostrate Spurge, Prostrate Vervain, Puncturevine, Purslane-Common, Ragweed, Red Sorrel, Redroot Pigweed, Redstem Filaree, Rough Cinquefoil, Rough Fleabane, Russian Pigweed, Russian Thistle, Scarlet Pimpernel, Scotch Thistle, Sheep Sorrel, Shepherds-purse, Slender Plantain, Smallflower Galinsoga, Smooth Dock, Smooth Pigweed, Sorrel, Sowthistle, Spanishneedles, Speedwell, Spiny Amaranth, Spiny Cocklebur, Spiny Sowthistle, Spotted Catsear, Spotted Spurge, Spurweed, Stinging Nettle, Strawberry-India Mock, Tall Nettle, Tall Vervain, Tansy Ragwort, Tansy Mustard, Thistle, Trailing Crownvetch, Tumble Mustard, Tumble Pigweed, Velvetleaf, Venice Mallow, Virginia Buttonweed, Virginia Creeper, Virginia Pepperweed, Wayleaf Bulbthistle, Western Clematis, Western Salsify, White Mustard, Wild Aster, Wild Buckwheat, Wild Carrot, Wild Four-O'clock, Wild Garlic, Wild Geranium, Wild Lettuce, Wild Marigold, Wild Mustard, Wild Onion, Wild Parsnip, Wild Radish, Wild Rape, Wild Strawberry, Wild Sweet Potato, Wild Vetch, Woodsorrel, Woolly Croton, Woolly Morningglory, Woolly Plantain, Wormseed, Yarrow, Yellow Rocket, Yellowflower Pepperweed.

ADVISORY FOR FERTILIZERS CONTAINING IRON

Keep this product off sidewalks, patios, driveways, etc. This fertilizer may contain iron which will stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

This product has been registered with the Washington State Department of Agriculture. When applied as directed, this fertilizer meets the Washington standards for arsenic, cadmium, cobalt, mercury, molybdenum, lead, nickel, selenium, and zinc. You have the right to receive specific information about Washington standards from the distributor of this product. Information received by the Washington State Department of Agriculture regarding the components in this product is available on the internet at <http://www.wa.gov/agr/>.

18-0-6

GUARANTEED ANALYSIS

Total Nitrogen (N)	18%
8.5% Ammoniacal Nitrogen	
9.5% Urea Nitrogen*	
Soluble Potash (K ₂ O)	6%
Boron (B)	0.02%
Copper (Cu)	0.05%
0.05% Water Soluble Copper (Cu)	
Iron (Fe)	5.00%
0.1% Water Soluble Iron (Fe)	
Manganese (Mn)	0.05%
0.05% Water Soluble Manganese (Mn)	
Zinc (Zn)	0.05%
0.05% Water Soluble Zinc (Zn)	

Derived From: Ammonium Phosphate, Polymer Coated Sulfur Coated Urea, Muriate of Potash, Sodium Borate, Copper Sulfate, Ferrous Sulfate, Iron Oxide, Manganese Sulfate, Zinc Sulfate.

*1.2% Slowly Available Nitrogen from Polymer Coated Sulfur Coated Urea.

10930-0515-TP

MANUFACTURED BY



230 FM 87 • BONHAM, TEXAS 75418

EPA Reg. No. 2217-883-7401 • EPA Est. No. 7401-TX-01

Visit Us At: www.fertilome.com

Product Questions? 855-270-4776

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

